







# IMAGE EVALUATION TEST TARGET (MT-3)



APPLIED IMAGE, Inc  
1653 East Main Street  
Rochester, NY 14609 USA  
Phone: 716/482-0300  
Fax: 716/288-5989

© 1993, Applied Image, Inc., All Rights Reserved



**CIHM  
Microfiche  
Series  
(Monographs)**

**ICMH  
Collection de  
microfiches  
(monographies)**



**Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques**

**© 1993**

Technical and Bibliographic Notes / Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

- ☐ Coloured covers/  
Couverture de couleur
- ☐ Covers damaged/  
Couverture endommagée
- ☐ Covers restored and/or laminated/  
Couverture restaurée et/ou pelliculée
- ☐ Cover title missing/  
Le titre de couverture manque
- ☐ Coloured maps/  
Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black)/  
Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations/  
Planches et/ou illustrations en couleur
- ☒ Bound with other material/  
Relié avec d'autres documents
- ☒ Tight binding may cause shadows or distortion  
along interior margin/  
La reliure serrée peut causer de l'ombre ou de la  
distorsion le long de la marge intérieure
- ☐ Blank leaves added during restoration may appear  
within the text. Whenever possible, these have  
been omitted from filming/  
Il se peut que certaines pages blanches ajoutées  
lors d'une restauration apparaissent dans le texte,  
mais, lorsque cela était possible, ces pages n'ont  
pas été filmées.

- ☐ Additional comments:/  
Commentaires supplémentaires:

This item is filmed at the reduction ratio checked below/  
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
						<input checked="" type="checkbox"/>					

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages/  
Pages de couleur
- ☐ Pages damaged/  
Pages endommagées
- ☐ Pages restored and/or laminated/  
Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed/  
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/  
Pages détachées
- ☒ Showthrough/  
Transparence
- ☒ Quality of print varies/  
Qualité inégale de l'impression
- ☐ Continuous pagination/  
Pagination continue
- ☐ Includes index(es)/  
Comprend un (des) index
- Title on header taken from:/  
Le titre de l'en-tête provient:
- ☐ Title page of issue/  
Page de titre de la livraison
- ☐ Caption of issue/  
Titre de départ de la livraison
- ☐ Masthead/  
Générique (périodiques) de la livraison

The copy filmed here has been reproduced thanks to the generosity of:




Library  
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol — (meaning "CONTINUED"), or the symbol ▽ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:



L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque  
Agriculture Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole — signifie "A SUIVRE", le symbole ▽ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

1894  
Exp. Farm Notes 2

EXPERIMENTAL FARM NOTES

No. 2.

---

CENTRAL EXPERIMENTAL FARM

DIVISION OF BOTANY,

DEPARTMENT OF AGRICULTURE.

---

POTATO BLIGHTS.

---

PUBLISHED BY DIRECTION OF THE HON. A. R. ANGERS, MINISTER OF AGRICULTURE.

OTTAWA  
GOVERNMENT PRINTING BUREAU  
1894.

EX  
CENT

FO

B

There are fe  
of more loss to  
have been aptly  
**EARLY BLIGHT**  
confounded und  
and "Potato ru  
similar in gener  
the attacks of t

(K

# EXPERIMENTAL FARM NOTES

## CENTRAL EXPERIMENTAL FARM.

(Department of Agriculture.)

### DIVISION OF BOTANY.

## POTATO BLIGHTS.

By JAMES FLETCHER, F.R.S.C., F.L.S.

There are few diseases of field crops which are the direct cause of more loss to the farmers of Canada than the two blights which have been aptly termed by Prof. L. R. Jones, of Vermont, the **EARLY BLIGHT** and **LATE BLIGHT** of potatoes. These are usually confounded under the various names "Potato rot," "Potato blight" and "Potato rust;" but, as a matter of fact, although somewhat similar in general appearance they are very distinct and are due to the attacks of two different vegetable parasites.



FIG. I.—THE EARLY BLIGHT.

(Kindly lent by Prof. L. R. Jones)

4

1. THE EARLY BLIGHT.—This disease is caused by the fungus *Macrosporium solani*, E. & M., and shows itself during the months of June and July, when greyish brown spots appear upon the older leaves. These soon become dry and crisp, and in bad cases the whole leaf is affected, so that nothing is left but the stems and the tubers stop growing.

The appearance of this disease is well shown in fig. 1.



FIG. 2.—THE LATE BLIGHT.  
(Kindly lent by Prof. L. R. Jones).

2. THE LATE BLIGHT, POTATO ROT.—This disease of the potato is due to the attack of a parasitic fungus, known by the name of *Phytophthora infestans*, D. By. The life history of this enemy is briefly as follows: The fungus passes the winter inside the potato tuber and is planted with it in the spring. As soon as the potato throws out its shoots, the parasite grows with it, running up through the tissues of the stems, and from about the end of July produces beneath the leaves an abundance of spores, or seed-like bodies. These are exceedingly minute, but are produced in such numbers that they frequently give a frost-like appearance to the under sides of the leaves. When these spores are produced on the leaves the appearance known as "rust" shows itself in the shape of dark brown spots, as shown in fig. 2, which are caused by the drying up of the tissues, from the parasite having used up their contents. From the rust stage all future infection takes place. Some of the spores are carried by the wind and falling upon the leaves of other adjacent plants, produce more rust spots, while

others fall reaching the ground as seen in the illustration. It is best known on the tubers, and in winter the diseased tubers are found.

In this stage about the first of the disease is present in a few leaves. It spreads rapidly of infection in a few days, and a

Careful examination of the tops five or six days early in July, will reveal the disease, for the first signs appear, for the first time, known to be in a large

Copied from the  
Linen  
Wash

To make a  
(blue vitriol)  
salt bag answer  
top of a coal  
that the bag  
the copper su  
vessel slake  
thin whitewash  
remove all l  
pour the lime  
the time. No  
mixture is rea

others falling to the ground are washed beneath the surface, and reaching the forming tubers produce the rot stage. The wet-rot, as seen in autumn in the tubers, is the form of this disease which is best known, but Potato-rot is really a dry-rot which kills the tubers, and in autumn the wet-rot follows as a result of decay. In winter the disease occurs in the tubers, as patches of hard, whitish, diseased tissue.

In this district the rust stage does not generally appear until about the first of August and this is the first evidence that blight is present in the field. As a rule the dark spots appear only on a few leaves at first, but if the weather be favourable the disease spreads rapidly from spores carried by the wind from these centres of infection, so that a large field may become diseased in a few days, and as a result the crop of potatoes be ruined.

#### REMEDY.

Careful experiments have shown that by spraying the potato tops five or six times at intervals of about two weeks, beginning early in July, for the early blight, and at the time the rust first appears, for the potato-rot, with the mixture of sulphate of copper and lime, known as the Bordeaux mixture, both of these injurious diseases can in a large measure be controlled.

#### BORDEAUX MIXTURE.

Copper sulphate .....	6 pounds.
Lime, fresh .....	4 "
Water .....	45 gallons.

*To make Bordeaux mixture*, take 6 pounds of copper sulphate (blue vitriol) and tie it up in a piece of thin cloth—an ordinary salt bag answers well—then suspend it from a stick laid across the top of a coal oil or other barrel half filled with clean water, so that the bag may be just beneath the surface of the water, when the copper sulphate will dissolve in an hour or two. In another vessel slake 4 pounds of fresh lime in sufficient water to make a thin whitewash. Strain this through a fine sieve or sack to remove all lumps. When the copper sulphate is all dissolved, pour the lime wash into the barrel slowly, stirring the mixture all the time. Now fill up the barrel to the top with water, and the mixture is ready for use.

*To apply this mixture* to the foliage, undoubtedly the best and cheapest way is to use a proper spraying pump and nozzle, but if these are not on hand, good results which will well repay the trouble, may be obtained by applying the mixture with watering cans supplied with fine roses. There are several different kinds of spraying pumps in the market; perhaps the most convenient for this work is a force pump attached to a barrel on wheels, to be drawn through the field by a horse. Smaller machines, known as Knapsack Sprayers, consist of a reservoir containing a small force pump, which can be carried upon a man's back. Both of these kinds of pumps can be purchased for about \$10 to \$20, and are now for sale by most of our seedsmen. It will be necessary to spray the field two or three times to protect the crop thoroughly. There is no danger of injuring the foliage with the above mixture.

A great advantage of this mixture is that Paris Green, the only practical remedy for the Colorado Potato-beetle, can be applied at the same time. To do this, mix from a quarter to half a pound of Paris Green with a little water so as to make a thick paste, and then add it to the 45 gallons of Bordeaux mixture; that is, it is used in exactly the same strength as with plain water.

These mixtures must be kept constantly stirred while being used, as both the lime in the Bordeaux mixture and the Paris Green sink quickly to the bottom of any mixture if left undisturbed.

*The time to apply.*—The Bordeaux mixture is a preventive remedy, and the time to apply it in any locality is just before the blights treated of usually appear there, the object being to keep the plants, during the whole of the time they are liable to injury, covered with the fungicidal preparation.

The early blight in this part of Canada generally appears at the end of June or early in July. The late blight or potato-rot seldom shows itself until August. Therefore, spraying should be begun early in July and repeated every two weeks at least until the end of August.

J. FLETCHER,

*Botanist.*

WM. SAUNDERS,

*Director.*

OTTAWA, July 1, 1894.

